

Hallux valgus and forefoot surgery

Hallux Valgus and Forefoot Surgery: An In-Depth Guide

Hallux valgus and forefoot surgery represent significant topics in foot and ankle medicine, particularly for individuals suffering from deformities that impair mobility, cause pain, and affect overall quality of life. Hallux valgus, commonly known as bunion, is one of the most prevalent forefoot deformities worldwide. When conservative treatments fail to alleviate symptoms, surgical intervention becomes necessary to restore function, reduce pain, and improve foot aesthetics.

Understanding the intricacies of hallux valgus and the various surgical options available is essential for patients, healthcare professionals, and podiatrists alike. This comprehensive guide aims to explore the causes, symptoms, diagnosis, and the latest advancements in forefoot surgery related to hallux valgus correction.

What Is Hallux Valgus?

Definition and Overview

Hallux valgus is a deformity characterized by lateral deviation of the great toe (hallux) and medial deviation of the first metatarsal bone. This results in a prominent bony bump on the inner side of the foot at the big toe's base. The deformity can vary in severity and may lead to discomfort, difficulty walking, and footwear limitations.

Causes of Hallux Valgus

Several factors contribute to the development of hallux valgus, including:

- Genetic predisposition: Family history increases risk.
- Footwear: Narrow, tight, or high-heeled shoes exert pressure on the forefoot.
- Foot biomechanics: Abnormal gait or abnormal pronation can stress the joint.
- Arthritis: Conditions like rheumatoid arthritis can accelerate deformity.
- Structural abnormalities: Flat feet or hypermobility contribute to deformity progression.
- Trauma: Previous injuries may predispose to bunion formation.

Symptoms and Signs

Patients with hallux valgus often present with:

- Visible bony bump on the inner side of the foot.
- Pain or tenderness around the bunion, especially with footwear.
- Swelling and redness.
- Restricted movement of the big toe.
- Corns or calluses due to pressure.
- Difficulty finding comfortable shoes.
- Changes in gait or walking pattern.

Diagnosing Hallux Valgus

Clinical Examination

Diagnosis begins with a detailed clinical assessment, including inspection and palpation of the foot. The healthcare provider will evaluate:

- The degree of deformity.
- Range of motion of the first MTP joint.
- Presence of tenderness, swelling, or skin changes.
- Gait analysis.

Imaging Studies

Radiographs are critical in confirming the diagnosis and planning treatment:

- Weight-bearing X-rays: Provide precise measurements of deformity.
- Key Angles Measured:
 - Hallux Valgus Angle (HVA): The angle between the longitudinal axes of the first metatarsal and the proximal phalanx.
 - Intermetatarsal Angle (IMA): The angle between the first and second metatarsal bones.

Normal values:

- HVA:
- IMA:

Deviations beyond these thresholds indicate severity and guide surgical planning.

Conservative Management of Hallux Valgus

Before considering surgery, non-invasive treatments are often attempted:

- Footwear modifications: Wide, low-heeled shoes.
- Padding and orthotics: To reduce pressure and improve biomechanics.
- NSAIDs: To control pain and inflammation.
- Physical therapy: Stretching and strengthening exercises.
- Footwear techniques: Taping or splinting to realign the toe temporarily.

While these measures can provide symptomatic relief, they do not correct the deformity. Persistent symptoms often necessitate surgical intervention.

Forefoot Surgery for Hallux Valgus: An Overview

Indications for Surgery

Surgical correction is typically indicated when:

- Pain persists despite conservative therapy.
- Deformity impairs daily activities.
- Progressive deformity threatens joint integrity.
- Significant functional impairment or gait disturbance.

Goals of Surgery

- Correct the alignment of the toe and metatarsal bones.
- Relieve pain.
- Restore normal foot biomechanics.
- Improve footwear options and aesthetics.

Types of Forefoot Surgical Procedures

Surgical options are tailored based on deformity severity, patient age, activity level, and bone quality. Common procedures include:

- Bunionectomy (Exostectomy)
- Osteotomy Procedures
- Arthrodesis

- Soft Tissue Procedures
- Combined Techniques

Each method has specific indications and advantages.

Common Surgical Techniques for Hallux Valgus

1. Chevron (Austin) Osteotomy

- A V-shaped cut performed at the distal first metatarsal.
- Allows lateral translation of the metatarsal head.
- Suitable for mild to moderate deformities.
- Often combined with soft tissue procedures.

2. Scarf Osteotomy

- A Z-shaped cut along the shaft of the first metatarsal.
- Provides greater stability and correction for moderate to severe deformities.
- Allows for rotation and lateral translation.
- Often used in combination with other procedures.

3. Reverdin Osteotomy

- Performed on the proximal phalanx.
- Corrects hallux valgus by realigning the toe.
- Usually combined with metatarsal osteotomies.

4. Proximal Osteotomies

- Performed at the base of the first metatarsal.
- Indicated for severe deformities.
- Examples include the Wilson or basal osteotomies.

5. Arthrodesis

- Fusion of the first MTP joint.

- Ideal for severe arthritis or rigid deformities.
- Provides pain relief and stability.

6. Soft Tissue Procedures

- Ligament and capsule releases or tightening.
- Tendon transfers to balance forces.
- Often combined with osteotomies for optimal correction.

Choosing the Right Surgical Approach

Factors Influencing Surgical Choice

- Severity of deformity (HVA and IMA angles).
- Age and activity level.
- Bone quality.
- Presence of arthritis.
- Patient expectations and cosmetic considerations.

Expected Outcomes

Most modern surgical techniques yield high success rates, with:

- Pain relief in over 90% of cases.
- Significant correction of deformity.
- Improved gait and footwear comfort.
- Low complication rates when performed by experienced surgeons.

Postoperative Care and Rehabilitation

Immediate Postoperative Period

- Use of a surgical shoe or boot.
- Elevation to reduce swelling.
- Limited weight-bearing protocols depending on the procedure.

Rehabilitation Timeline

- Weeks 1-6: Rest, immobilization, and gradual weight-bearing.
- Weeks 6-12: Gradual return to normal activities.
- Physical therapy: To restore strength and range of motion.

Potential Complications

While generally safe, forefoot surgery carries risks such as:

- Infection.
- Nerve injury.
- Recurrence of deformity.
- Overcorrection or undercorrection.
- Non-union or delayed union of osteotomies.
- Stiffness or residual deformity.

Advances in Forefoot Surgery

Minimally Invasive Techniques

Recent innovations include percutaneous osteotomies that reduce soft tissue disruption, minimize scarring, and promote faster recovery.

Computer-Assisted Surgery

Preoperative planning with 3D imaging and intraoperative navigation enhance precision and outcomes.

Biological Enhancements

Use of bone grafts, growth factors, and scaffolds to improve healing, especially in complex cases.

Conclusion

Hallux valgus and forefoot surgery continue to evolve with advancements in surgical techniques, imaging, and postoperative management. A personalized approach considering deformity severity, patient needs, and surgeon expertise ensures optimal outcomes. When conservative measures fail, surgical correction offers a reliable solution to restore foot function, alleviate pain, and improve quality of life.

Patients considering surgery should consult with a specialist to discuss the most appropriate

procedure, understand potential risks, and set realistic expectations for recovery and long-term results. With proper planning and execution, forefoot surgeries can significantly enhance mobility and comfort for those affected by hallux valgus.

Hallux valgus and forefoot surgery are topics of significant interest within the field of foot and ankle orthopedics, given their prevalence and impact on patient quality of life. Hallux valgus, commonly known as a bunion, is a deformity characterized by lateral deviation of the big toe and medial prominence of the first metatarsal head. When conservative measures fail to alleviate symptoms or deformity progresses, surgical intervention becomes necessary. Forefoot surgery encompasses a broad spectrum of procedures designed to correct deformities, relieve pain, and restore function.

In this comprehensive guide, we will explore the anatomy and pathology of hallux valgus, indications for surgery, various surgical techniques, postoperative care, and emerging trends in forefoot correction. Whether you are a healthcare professional, a student, or a patient seeking in-depth understanding, this article aims to provide clarity on this complex yet common condition.

Understanding Hallux Valgus: Anatomy and Pathophysiology

Anatomy of the Forefoot

The forefoot comprises five metatarsal bones and the proximal phalanges of the toes. The first metatarsal is the largest and most medial, articulating with the medial cuneiform. The big toe (hallux) is aligned in an anatomically straight position but can deviate laterally in hallux valgus.

Pathophysiology of Hallux Valgus

Hallux valgus involves a complex interplay of anatomical, biomechanical, and genetic factors:

- Bony deformity: Lateral deviation of the proximal phalanx and medial deviation of the first metatarsal.
- Soft tissue imbalance: Laxity of the medial capsule, medial collateral ligament, adductor hallucis muscle, and the plantar fascia.
- Biomechanical factors: Overpronation, abnormal gait, footwear choices (narrow, high-heeled shoes).
- Genetics: Family history increases risk.

As the deformity progresses, the medial eminence (bunion) enlarges, and the joint becomes unstable, often leading to pain, bursitis, and difficulty wearing shoes.

Indications for Forefoot Surgery in Hallux Valgus

Surgical correction is typically considered when:

- Persistent pain despite conservative measures such as footwear modification, orthoses, or medication.
- Progressive deformity leading to functional impairment.
- Cosmetic concerns affecting patient satisfaction.
- Decreased quality of life due to gait abnormalities or discomfort.

Key considerations before surgery include:

- Degree of deformity (measured by angles such as hallux valgus angle (HVA) and intermetatarsal angle (IMA))
- Patient's age, activity level, and overall health
- Presence of arthritis or joint degeneration
- Patient expectations and compliance

Surgical Techniques for Hallux Valgus Correction

Forefoot surgery for hallux valgus is diverse, with procedures tailored to the severity and specific deformity components. The overarching goal is to realign the toe, stabilize the joint, and relieve symptoms.

Common Surgical Procedures

- Soft Tissue Procedures
 - Medial capsule reefing: Tightening lax soft tissues.
 - Adductor hallucis release: Reducing lateral pull on the big toe.
 - Bursotomy: Removing inflamed bursa over the bunion.
- Bony Osteotomies

Osteotomies are cuts made in the bone to correct deformity:

- Chevron (Austin) Osteotomy: The most common for mild to moderate deformities; involves a

V-shaped cut in the distal first metatarsal.

- Reverse Chevron (Wilson) Osteotomy: Similar to Chevron, but with modifications for specific deformities.
- Distal Soft Tissue Procedure (Lapidus): Fusion of the first tarsometatarsal joint, used for severe or hypermobile deformities.
- Proximal Osteotomies: Such as the proximal chevron or proximal metatarsal osteotomy, used for more severe deformities.
- Arthrodesis
- Fusion of the first metatarsophalangeal joint; indicated in arthritic cases or severe deformities.
- Other Techniques
- Osteotomy with implant fixation: Enhanced stability using screws, plates.
- Minimally invasive surgery: Percutaneous techniques gaining popularity for less soft tissue disruption.

Step-by-Step Overview of Common Procedures

A. Chevron (Austin) Osteotomy

- Performed through a lateral incision over the first metatarsal head.
- A V-shaped cut is made approximately 1.5 to 2 cm distal to the metatarsal head.
- The distal fragment is shifted medially to correct the IMA.
- Fixation with screws or wires stabilizes the osteotomy.
- Soft tissue balancing complements the bony correction.

B. Lapidus Procedure

- Involves fusion of the first tarsometatarsal joint.
- Suitable for hypermobile or severe deformities.
- Provides robust correction and stability.
- Typically involves placement of screws or plates to maintain fusion.

Postoperative Care and Rehabilitation

Proper postoperative management is critical to ensure healing and optimal outcomes.

Initial Phase (0-2 weeks):

- Immobilization in a postoperative shoe or cast.
- Non-weight bearing or partial weight-bearing as advised.
- Pain management and edema control.

Intermediate Phase (2-6 weeks):

- Transition to weight-bearing as tolerated.
- Wound care and monitoring for signs of infection.
- Initiation of gentle range-of-motion exercises.

Rehabilitation Phase (6+ weeks):

- Gradual return to normal footwear.
- Strengthening exercises.
- Gait training and physical therapy if needed.

Potential Complications:

- Infection
- Nonunion or delayed union
- Residual deformity or recurrence
- Overcorrection leading to hallux varus
- Neurovascular injury

Emerging Trends and Future Directions

Advances in forefoot surgery continue to evolve, with notable trends including:

- Minimally invasive techniques: Smaller incisions, reduced soft tissue trauma, faster recovery.
- Percutaneous osteotomies: Use of specialized tools for bone cuts through minimal incisions.

- Biological augmentation: Use of bone grafts or growth factors to enhance healing.
- 3D imaging and planning: Preoperative virtual simulations for precise correction.
- Patient-specific implants: Customized fixation devices for complex deformities.

Conclusion: Choosing the Right Surgical Approach

Successful correction of hallux valgus and forefoot deformities hinges on a comprehensive understanding of the deformity's components, patient-specific factors, and surgical expertise. A tailored approach ensures optimal alignment, function, and patient satisfaction.

Patients should be thoroughly counseled regarding expectations, risks, and the rehabilitation process. Collaboration between surgeons, radiologists, physiotherapists, and patients is essential for achieving the best outcomes.

As research progresses and techniques become less invasive, the future of forefoot surgery promises improved results with quicker recoveries and fewer complications. Whether through traditional osteotomies or emerging minimally invasive methods, the goal remains consistent: restoring a pain-free, functional, and cosmetically acceptable foot.

Note: For specific surgical planning or individualized recommendations, consultation with a qualified foot and ankle specialist is essential.

Question What are the common symptoms of hallux valgus that indicate the need for forefoot surgery? Common symptoms include a visible bunion, pain or tenderness around the big toe joint, swelling, limited toe movement, and difficulty wearing narrow shoes. Persistent discomfort despite conservative treatment may warrant surgery. What are the latest surgical techniques used to treat hallux valgus? Recent advancements include minimally invasive procedures such as percutaneous osteotomies, Lapidus procedures, and arthroscopic corrections, which aim to reduce recovery time and improve cosmetic outcomes. How do I know if I am a good candidate for forefoot surgery for hallux valgus? Candidates typically have significant bunion deformity causing pain or functional impairment, and have not responded to conservative treatments. A thorough evaluation by a foot and ankle specialist is essential to determine suitability. What is the recovery process like after hallux valgus surgery? Recovery generally involves a period of immobilization, weight-bearing restrictions, and physical therapy. Most patients can expect to return to normal activities within 6 to 8 weeks, with full recovery potentially taking several months. Are there any risks or complications associated with forefoot surgery for hallux valgus? Possible risks include infection, nerve injury, recurrence of deformity, stiffness, and issues related to hardware. Choosing an experienced surgeon and following postoperative care instructions can minimize these risks. What are the benefits of early surgical intervention for hallux valgus? Early surgery can alleviate pain, prevent progression of deformity, improve foot function, and reduce the likelihood of secondary issues such as arthritis or hammertoes. How does forefoot surgery impact long-term foot health and function?

Properly performed surgery can restore normal alignment, relieve pain, and improve gait. However, long-term success depends on patient health, adherence to postoperative protocols, and addressing any underlying biomechanical issues. What are the latest developments in postoperative care for hallux valgus surgery? Emerging approaches include enhanced rehabilitation protocols, use of advanced surgical fixation devices, and patient-specific orthoses to improve outcomes and reduce recovery times.

Related keywords: hallux valgus, bunion correction, forefoot osteotomy, bunionectomy, minimally invasive bunion surgery, first metatarsal deformity, toe alignment, bunion pain relief, forefoot reconstruction, postoperative rehabilitation

Operative techniques foot and ankle surgery

operative techniques foot and ankle surgery

Foot and ankle surgery encompasses a diverse array of operative techniques aimed at correcting deformities, repairing injuries, and restoring function to the lower extremity. Advancements in surgical methods have significantly improved patient outcomes, reduced recovery times, and minimized complications. This comprehensive guide explores the most common and effective operative techniques used in foot and ankle surgery, providing valuable insights for orthopedic surgeons, residents, and healthcare professionals involved in limb preservation and rehabilitation.

Overview of Foot and Ankle Surgical Interventions

Foot and ankle surgeries address a wide spectrum of conditions including fractures, ligament injuries, tendinopathies, deformities such as hallux valgus, and degenerative diseases like arthritis. The choice of operative technique depends on factors like the specific pathology, patient health status, and functional demands.

Common Foot and Ankle Surgical Procedures

1. Arthroscopic Techniques

Arthroscopy provides minimally invasive access to the joint space, enabling diagnosis and treatment with reduced soft tissue trauma.

- Ankle Arthroscopy: Used for osteochondral lesions, synovitis, loose body removal, and joint

stabilization.

- Subtalar and Midtarsal Arthroscopy: Facilitates assessment and management of hindfoot deformities and coalition resections.

Advantages:

- Reduced postoperative pain
- Shorter hospital stays
- Quicker return to activity

2. Osteotomies

Osteotomies involve cutting and realigning bones to correct deformities such as hallux valgus or cavus foot.

- Austin (Chevron) Osteotomy: For mild to moderate hallux valgus correction.
- Scarf Osteotomy: Allows greater deformity correction with increased stability.
- Dwyer Osteotomy: Used for flexible flatfoot deformity.

Key Points:

- Precise angular correction is critical.
- Fixation devices like screws or plates ensure stability.

3. Arthrodesis (Joint Fusion)

A definitive procedure for stabilizing painful or arthritic joints.

- Tibiotalar Arthrodesis: Fusion of the ankle joint.
- Subtalar Arthrodesis: Stabilization of the subtalar joint.
- First MTP Arthrodesis: For severe hallux rigidus.

Technique:

- Precise joint preparation and removal of cartilage.
- Compression fixation with screws or plates.

- Postoperative immobilization for fusion success.

4. Soft Tissue Repairs and Reconstructions

Address ligament injuries, tendinopathies, or soft tissue defects.

- Lateral Ligament Repair: For ankle sprains, often using suture anchors.
- Achilles Tendon Repair: Open or minimally invasive techniques for ruptures.
- Plantar Fascia Release: For recalcitrant plantar fasciitis.

Considerations:

- Proper tensioning is essential.
- Postoperative immobilization may be required.

5. Fracture Fixation Techniques

Management of foot and ankle fractures is critical for restoring anatomy and function.

- Open Reduction and Internal Fixation (ORIF): For displaced fractures.
- Percutaneous Fixation: Minimally invasive approach for select fractures.
- External Fixation: For complex or open fractures.

Materials Used:

- Screws
- Plates
- Pins
- External fixators

6. Tendon Transfers and Reconstructive Procedures

Used in cases of tendon deficiency or dysfunction.

- Posterior Tibial Tendon Transfer: To correct flatfoot deformity.
- Peroneal Tendon Transfer: For eversion weakness.

- Flexor Tendon Transfers: In cases of nerve injury.

Emerging and Advanced Techniques

1. Minimally Invasive and Percutaneous Techniques

Advances in imaging and instrumentation have enabled less invasive procedures with comparable outcomes.

- Percutaneous osteotomies for bunion correction.
- Arthroscopic-assisted ligament repairs.

2. 3D Printing and Custom Implants

Customized implants and guides improve anatomical accuracy, especially in complex deformity corrections.

3. Biological Enhancements

Use of growth factors, stem cells, and bone grafts to enhance healing in non-union or osteoarthritic cases.

Preoperative Planning and Intraoperative Considerations

Effective foot and ankle surgery relies heavily on meticulous planning.

Preoperative Steps:

- Detailed clinical examination.
- Imaging studies: X-rays, CT scans, MRI.
- Assessment of patient-specific factors like vascular status and comorbidities.

Intraoperative Principles:

- Adequate exposure while minimizing soft tissue damage.
- Precise osteotomy angles and fixation.
- Restoration of normal anatomy and biomechanics.
- Ensuring stable fixation for optimal healing.

Postoperative Care and Rehabilitation

Successful outcomes depend on structured postoperative protocols:

- Immobilization with casts or braces.
- Gradual weight-bearing as per fixation stability.
- Physical therapy focusing on range of motion, strength, and proprioception.
- Monitoring for complications such as infection, non-union, or hardware failure.

Potential Complications and Their Management

- Infection: Early detection and antibiotic therapy.
- Non-union or Malunion: May require revision surgery.
- Neurovascular Injury: Careful dissection and intraoperative nerve monitoring.
- Hardware-related issues: Removal or revision as necessary.

Conclusion

Operative techniques in foot and ankle surgery are continuously evolving, driven by technological innovations and a deeper understanding of biomechanics. Surgeons must tailor interventions to individual patient needs, employing minimally invasive methods when appropriate, while ensuring anatomical restoration and functional recovery. Mastery of these operative techniques, combined with meticulous preoperative planning and postoperative management, is essential for achieving optimal patient outcomes in foot and ankle surgical care.

Keywords: foot surgery, ankle surgery, osteotomy, arthrodesis, minimally invasive, fracture fixation, ligament repair, tendinopathy, deformity correction, surgical techniques

Operative Techniques Foot and Ankle Surgery

Foot and ankle surgery encompasses a diverse array of operative procedures aimed at addressing a multitude of pathologies affecting these complex and biomechanically critical parts of the human body. As the foundation of mobility, stability, and weight-bearing, the foot and ankle demand precise surgical intervention when conservative measures fail. The evolution of operative techniques in foot and ankle surgery has been driven by advances in imaging, fixation devices, minimally invasive approaches, and a deeper understanding of biomechanics. This article provides a comprehensive overview of current operative techniques, emphasizing indications, procedural details, advantages, limitations, and emerging trends.

Overview of Foot and Ankle Surgical Indications

Foot and ankle surgeries are indicated for a broad spectrum of conditions including traumatic injuries, degenerative diseases, deformities, infections, and tumors. Common indications include:

- Fractures and dislocations
- Ligamentous injuries, such as ankle sprains and chronic instability
- Tendinopathies and ruptures
- Arthritic changes, such as osteoarthritis and rheumatoid arthritis
- Deformities like hallux valgus, cavus foot, flatfoot, and clubfoot
- Neuropathic conditions, including Charcot neuroarthropathy
- Infections and neoplastic lesions

Understanding the specific pathology guides the choice of operative technique, which ranges from minimally invasive procedures to complex reconstructions.

Trauma Management in Foot and Ankle Surgery

Trauma constitutes a significant portion of foot and ankle surgeries, often requiring precise reduction and stable fixation to restore function.

Open Reduction and Internal Fixation (ORIF)

Procedure Overview:

ORIF remains the gold standard for displaced fractures of the calcaneus, talus, distal tibia, and other bones. It involves surgical exposure, reduction of fracture fragments, and stabilization with plates, screws, or nails.

Key Techniques:

- Lateral approach for calcaneal fractures
- Anterior or posterior approaches for talar neck fractures
- Use of locking plates and screws for stable fixation

Pros:

- Precise anatomical reduction

- Early mobilization possibilities
- High union rates

Cons:

- Soft tissue complications
- Infection risk
- Longer operative times

Minimally Invasive and Percutaneous Fixation

With technological advances, percutaneous techniques are increasingly employed for specific fractures, offering reduced soft tissue disruption.

Features:

- Smaller incisions
- Reduced wound complications
- Faster recovery

Limitations:

- Limited visualization
- Potential for suboptimal reduction in complex fractures

Deformity Correction Techniques

Correcting deformities is pivotal in restoring biomechanics and preventing degenerative changes.

Hallux Valgus (Bunion) Surgery

Common Procedures:

- Distal osteotomies (e.g., Chevron osteotomy)
- Proximal osteotomies (e.g., Wilson osteotomy)
- Arthrodesis for severe deformities

Operative Techniques:

- Lateral soft tissue release
- Osteotomy fixation with screws or plates
- Realignment of the first metatarsal

Pros:

- Symptom relief
- Improved foot alignment
- Durable correction

Cons:

- Recurrence if improperly performed
- Potential for stiffness or transfer metatarsalgia

Flatfoot and Cavus Foot Corrections

Procedures include medializing calcaneal osteotomy, lateral column lengthening, and flexor tendon transfers.

Features:

- Realignment of the medial arch
- Restoring normal biomechanics

Pros:

- Pain relief
- Improved gait

Cons:

- Complex procedures

- Risk of overcorrection or undercorrection

Arthrodesis and Joint Preservation

Arthrodesis, or joint fusion, is frequently employed for end-stage arthritis or instability.

Foot and Ankle Arthrodesis Techniques

Types:

- Ankle fusion (tibiotalar joint)
- Subtalar fusion
- Midfoot (naviculocuneiform, Lisfranc) fusion

Procedural Approach:

- Rigid fixation with screws, plates, or bioabsorbable devices
- Proper positioning to maintain biomechanics

Advantages:

- Pain relief
- Durable stability

Disadvantages:

- Loss of joint mobility
- Adjacent joint degeneration over time

Joint Preservation Procedures

In early arthritis, procedures such as cheilectomy, osteotomy, or joint debridement aim to delay fusion.

Features:

- Maintains joint motion
- Less invasive

Limitations:

- May not be suitable for advanced disease
- Variable long-term outcomes

Soft Tissue Reconstruction Techniques

Reconstructive procedures are vital in chronic instability, tendon ruptures, or neurogenic conditions.

Ligament Repair and Reconstruction

Procedures:

- Broström procedure for lateral ankle ligament repair
- Tendon rerouting or augmentation techniques

Features:

- Restores stability
- Often combined with osteotomies

Pros:

- Preservation of native tissues
- Good functional outcomes

Cons:

- Risk of re-injury
- Technical demands

Achilles Tendon Repair

Techniques include open, minimally invasive, or percutaneous repair, often using suture anchors or core sutures.

Advantages:

- Restores plantarflexion strength
- Reduces re-rupture rates

Disadvantages:

- Wound healing issues in open repair
- Re-rupture risk if not properly rehabilitated

Innovations in Foot and Ankle Surgery

Recent advancements have revolutionized operative techniques, making procedures safer and more effective.

Minimally Invasive and Arthroscopic Techniques

Features:

- Reduced soft tissue trauma
- Faster recovery
- Enhanced visualization for joint pathology

Applications:

- Arthroscopic ankle arthrodesis
- Subtalar joint procedures
- Debridement of osteochondral lesions

Pros:

- Less postoperative pain
- Decreased wound complications

Cons:

- Steep learning curve
- Equipment costs

Computer-Assisted and Navigation-Guided Surgery

These technologies improve accuracy for complex reconstructions and deformity corrections.

Features:

- 3D planning
- Real-time navigation

Advantages:

- Precise alignment
- Potentially better outcomes

Limitations:

- Increased operative time
- Need for specialized training

Postoperative Care and Rehabilitation

Successful surgical outcomes depend not only on the operative technique but also on meticulous postoperative management.

- Early mobilization protocols versus immobilization
- Weight-bearing decisions based on fixation stability
- Physical therapy to regain strength and function

Emerging trends include the integration of regenerative medicine, such as platelet-rich plasma and stem cell therapies, to enhance healing, though their roles in operative techniques are still under investigation.

Conclusion

Operative techniques in foot and ankle surgery are diverse and continually evolving, driven by technological innovations and a deeper understanding of biomechanics. The choice of procedure depends on the pathology, patient factors, and surgeon expertise. While minimally invasive methods and advanced fixation devices offer significant benefits, they require specialized training and careful patient selection. As research progresses, future developments promise even less invasive, more precise, and biologically friendly approaches, ultimately improving patient outcomes and quality of life.

In summary:

- A tailored approach considering individual pathology and patient needs is essential.
- Advances in imaging, fixation, and minimally invasive techniques have expanded surgical options.
- Understanding the pros and cons of each technique allows for optimal decision-making.
- Multidisciplinary care and rehabilitation are integral to success.

Foot and ankle surgery remains a dynamic field, with ongoing innovations poised to further enhance operative outcomes for patients worldwide.

Question What are the key considerations when choosing an operative technique for hallux valgus correction? Selecting the appropriate operative technique depends on the severity of deformity, the presence of degenerative changes, patient comorbidities, and surgeon experience. Common options include osteotomies like Chevron or scarf osteotomies, with the goal of realigning the first metatarsal and correcting soft tissue imbalance for optimal functional and cosmetic outcomes. How does minimally invasive foot and ankle surgery compare to traditional open procedures? Minimally invasive techniques often result in smaller incisions, reduced soft tissue trauma, faster recovery times, and decreased postoperative pain. However, they require specialized training and instrumentation, and their suitability depends on the specific pathology and deformity severity. Proper patient selection is crucial for optimal outcomes. What are the common fixation methods used in foot and ankle osteotomies? Fixation methods include screws (lag screws, cortical or cancellous), plates and screws, Kirschner wires (K-wires), and bioabsorbable materials. The choice depends on the osteotomy type, stability requirements, bone quality, and surgeon preference to ensure proper healing and deformity correction. What are the indications for arthroscopic versus open repair in ankle ligament injuries? Arthroscopic repair is indicated for specific ligament injuries, such as anterior talofibular ligament tears, especially when concomitant intra-articular pathology exists. Open repair may be preferred for complex, chronic, or multi-ligament injuries, or when extensive debridement and reconstruction are necessary to restore stability. How do you minimize complications during foot and ankle reconstructive surgery? Complication prevention involves

meticulous surgical planning, gentle tissue handling, accurate anatomical reduction, stable fixation, and careful soft tissue management. Preoperative imaging, intraoperative fluoroscopy, and adherence to aseptic techniques further reduce risks such as infection, nerve injury, or nonunion. What advances have improved nerve decompression techniques in foot and ankle surgery? Advances include the use of high-resolution ultrasound for diagnosis, minimally invasive decompression approaches, and intraoperative nerve monitoring. These innovations enhance precision, reduce tissue trauma, and improve outcomes in conditions like tarsal tunnel syndrome. When is a tendon transfer indicated in foot and ankle operative procedures? Tendon transfers are indicated in cases of muscular weakness, nerve palsies, or deformities such as drop foot or cavovarus deformity. They help restore function, improve biomechanics, and correct deformities by redirecting or augmenting the action of specific muscle groups. What are the considerations for choosing between internal and external fixation in ankle fractures? Internal fixation is preferred for most stable fractures requiring precise anatomical reduction, providing rigid stabilization. External fixation may be indicated in open fractures, severe soft tissue injury, or when internal fixation is contraindicated due to infection risk or poor soft tissue condition. The choice depends on fracture pattern, soft tissue status, and patient factors.

Related keywords: foot and ankle surgery, surgical techniques, minimally invasive surgery, ligament repair, tendon reconstruction, ankle arthroscopy, fracture fixation, ankle stabilization, arthrodesis, osteotomy

Mann s surgery of the foot and ankle expert consu

mann s surgery of the foot and ankle expert consu is a leading specialty that focuses on diagnosing, treating, and rehabilitating complex issues related to the foot and ankle. For individuals experiencing persistent pain, deformities, or injuries in these crucial parts of the body, consulting an experienced foot and ankle surgeon is essential. Such specialists possess the advanced skills and knowledge needed to perform intricate surgeries, whether for traumatic injuries, deformities, or degenerative conditions. This article explores the scope of Mann's surgery of the foot and ankle, the expertise involved, common procedures, and why choosing the right specialist can significantly impact outcomes.

Understanding Mann's Surgery of the Foot and Ankle

What Is Mann's Surgery?

Mann's surgery refers broadly to surgical interventions performed by foot and ankle specialists to address a variety of conditions affecting the foot, ankle, and lower leg. Named after pioneering

surgeons or institutions, this term emphasizes advanced surgical techniques tailored to restore function, relieve pain, and improve quality of life. While not a single procedure, it encompasses a range of surgical treatments designed for complex or severe cases that cannot be managed conservatively.

The Role of a Foot and Ankle Expert

A foot and ankle expert is typically an orthopedic surgeon with specialized training in the musculoskeletal system of the lower extremities. These professionals have undergone extensive education and hands-on training in procedures such as deformity correction, joint replacement, fracture fixation, and soft tissue reconstruction. Their expertise ensures that patients receive personalized care that considers both the mechanical and biological aspects of foot and ankle health.

Common Conditions Treated with Mann's Surgery

Traumatic Injuries

Injuries such as fractures, dislocations, and ligament tears are common reasons for surgical intervention. These injuries often require precise realignment and stabilization to restore normal function.

Deformities and Congenital Conditions

Conditions like bunions, hammertoes, flat feet, high arches, and clubfoot often necessitate corrective surgery to improve biomechanics and alleviate discomfort.

Degenerative Diseases

Osteoarthritis and rheumatoid arthritis can severely impair mobility. Surgical options such as joint fusion or replacement help reduce pain and preserve joint function.

Soft Tissue Pathologies

Tendinitis, ligament tears, and tumors may require surgical repair or excision, especially when conservative treatments fail.

Types of Surgical Procedures Performed

Common Foot and Ankle Surgeries

- Bunionectomy (Hallux Valgus Correction)

A procedure to realign the big toe joint, relieving pain and deformity.

- Hammertoe Correction

Surgical straightening of toes affected by abnormal bending.

- Ankle Arthroscopy

A minimally invasive technique for diagnosing and treating joint issues such as cartilage damage or loose bodies.

- Achilles Tendon Repair

Surgical repair of torn or ruptured Achilles tendons to restore strength and mobility.

- Arthrodesis (Joint Fusion)

Joining bones in the foot or ankle to eliminate painful movement, often used in severe arthritis cases.

- Total Ankle Replacement

Replacing the damaged ankle joint with a prosthesis to preserve motion and reduce pain.

- Fracture Fixation

Using plates, screws, and pins to stabilize broken bones for proper healing.

The Surgical Process: What Patients Can Expect

Preoperative Evaluation

- Detailed medical history and physical examination.
- Imaging studies such as X-rays, MRI, or CT scans.
- Discussing surgical options, risks, and expected outcomes.

Surgical Planning

- Customized surgical approach based on the specific condition.
- Consideration of patient's overall health, activity level, and goals.

The Surgery

- Performed under local, regional, or general anesthesia.
- Use of advanced techniques to ensure minimal tissue trauma.
- Intraoperative imaging to enhance precision.

Postoperative Care

- Pain management strategies.
- Immobilization and weight-bearing restrictions.
- Physical therapy to regain strength and mobility.
- Regular follow-up appointments to monitor healing.

Benefits of Choosing a Mann's Surgery Expert

Specialized Skills and Experience

Expert surgeons have extensive experience with complex foot and ankle procedures, increasing the likelihood of successful outcomes.

Advanced Surgical Techniques

Access to the latest minimally invasive methods reduces recovery time, minimizes scarring, and lowers complication risks.

Personalized Treatment Plans

Experts tailor interventions to individual needs, considering age, activity level, and specific pathology.

Comprehensive Care

From diagnosis to rehabilitation, specialized centers offer integrated services, ensuring continuity of care.

Factors to Consider When Selecting a Foot and Ankle Surgeon

Credentials and Experience

- Board certification in orthopedic surgery with specialization in foot and ankle.
- Proven track record with similar cases.

Facilities and Technology

- Access to state-of-the-art surgical tools and imaging equipment.
- Accredited surgical centers.

Patient Reviews and Testimonials

- Feedback from previous patients about their experiences and outcomes.

Consultation and Communication

- Willingness to answer questions and discuss all options thoroughly.
- Clear explanation of risks, benefits, and recovery expectations.

Rehabilitation and Recovery

Post-Surgical Rehabilitation

Effective recovery involves a multidisciplinary approach combining physical therapy, pain management, and patient education.

Typical Timeline

- Immediate postoperative period: immobilization and limited weight-bearing.

- Short-term (2-6 weeks): gradual mobilization and physical therapy.
- Long-term (3-12 months): full activity resumption, sports, or work, depending on the procedure.

Importance of Adherence

Following surgeon instructions is critical to optimize healing, prevent complications, and restore function.

Conclusion: Why Expert Consultation Matters

Mann's surgery of the foot and ankle encompasses a wide array of complex procedures aimed at correcting deformities, repairing injuries, and restoring mobility. Consulting a specialized foot and ankle expert ensures that patients benefit from precise diagnosis, advanced surgical techniques, and comprehensive postoperative care. If you are experiencing chronic foot or ankle pain, deformity, or injury that impairs your daily activities, seeking expert consultation is a crucial step toward effective treatment and improved quality of life. Remember, the right surgical intervention by a skilled specialist can make all the difference in achieving lasting relief and optimal function.

Mann's Surgery of the Foot and Ankle: An Expert Review and Comprehensive Guide

When it comes to foot and ankle surgical interventions, Mann's surgery stands out as a specialized and highly regarded approach among orthopedic surgeons and podiatrists. Developed and refined over decades, this surgical technique offers targeted solutions for a variety of complex foot and ankle conditions. In this article, we delve deep into Mann's surgery, exploring its origins, indications, procedural steps, advantages, challenges, and expert opinions, providing a thorough resource for clinicians, patients, and healthcare enthusiasts alike.

Understanding Mann's Surgery of the Foot and Ankle

Mann's surgery is a specialized surgical intervention primarily designed to correct deformities, facilitate joint preservation, and improve function in various foot and ankle pathologies. It is not a single procedure but rather a set of techniques and principles that can be tailored to address specific issues such as arthritis, deformities, tendinopathies, and instability.

Historical Context and Development

Dr. Mann, a renowned orthopedic surgeon specializing in foot and ankle surgery, pioneered this approach in the late 20th century. His work was motivated by the need for minimally invasive, joint-preserving procedures that could restore biomechanics and reduce recovery time. Over the years, Mann's principles have been adapted and integrated into modern surgical practice, emphasizing patient-centered outcomes and functional restoration.

Core Principles of Mann's Surgery

Understanding the foundational concepts of Mann's surgery is essential to appreciate its versatility and effectiveness. Its core principles include:

- Joint Preservation

Rather than opting for complete joint replacement or fusion, Mann's surgery emphasizes preserving native joint structures whenever feasible. This approach maintains natural biomechanics and reduces the risk of adjacent joint degeneration.

- Anatomical Realignment

Restoring normal anatomical alignment is crucial. Mann's procedures focus on correcting deformities at their source—be it through osteotomies, soft tissue balancing, or ligament reconstruction—to ensure functional motion and stability.

- Minimally Invasive Techniques

Whenever possible, the approach favors minimally invasive methods to reduce soft tissue damage, minimize scarring, and accelerate recovery.

- Customized Treatment Plans

Recognizing patient variability, Mann's surgery is tailored to individual anatomy, pathology, and functional demands, often combining multiple techniques for optimal outcomes.

Common Conditions Treated with Mann's Surgery

Mann's surgical principles are applicable to a broad spectrum of foot and ankle disorders, including:

- Hallux Valgus (Bunion Deformity)
- Claw Toe and Hammer Toe Deformities
- Tarsal Coalition
- Flatfoot Deformity
- Ankle Osteoarthritis

- Tendon Pathologies (e.g., Achilles tendinopathy)
- Recurrent Ankle Instability

Each condition requires a specific adaptation of Mann's techniques, emphasizing the importance of expert assessment and planning.

Procedural Overview: Techniques and Approaches

While Mann's surgery comprises various procedures, several core techniques are frequently employed or adapted:

1. Osteotomies for Deformity Correction

Osteotomy involves precise cutting and realignment of bones to correct deformities like bunions or flatfoot. Mann's approach emphasizes minimally invasive osteotomies that preserve blood supply and soft tissue attachments.

- Examples include:
- Chevron or scarf osteotomies for hallux valgus
- Dwyer osteotomy for calcaneal deformities
- Medializing calcaneal osteotomy for hindfoot valgus

2. Soft Tissue Balancing

Correcting soft tissue imbalances is key to restoring function. Techniques involve:

- Ligament reconstructions
- Tendon transfers
- Capsular tightening or release

3. Arthrodesis and Arthroplasty

In cases of severe joint degeneration, Mann's principles support joint fusion (arthrodesis) for stability or joint-preserving procedures when possible.

4. Tendon and Ligament Reconstruction

Reconstructing or reinforcing soft tissues ensures joint stability and proper biomechanics, especially in cases of recurrent instability.

Advantages of Mann's Surgery

Patients and clinicians often favor Mann's techniques for their numerous benefits:

- **Preservation of Natural Joints:** Promotes better long-term function and reduces the need for prostheses.
- **Reduced Recovery Time:** Minimally invasive methods result in less postoperative pain and faster mobilization.
- **Customized Approach:** Tailored procedures lead to higher satisfaction and better functional outcomes.
- **Biomechanical Restoration:** Focus on anatomical realignment improves gait and reduces compensatory issues.
- **Lower Complication Rates:** Precise techniques minimize soft tissue trauma and related complications.

Challenges and Limitations

Despite its advantages, Mann's surgery is not without challenges:

- **Technical Complexity:** Requires specialized training and experience.
- **Patient Selection:** Not all conditions are suitable; advanced deformities may necessitate alternative procedures.
- **Long-Term Outcomes:** Continued research is needed to fully establish durability in some cases.
- **Learning Curve:** Surgeons must develop proficiency to optimize results and minimize complications.

Expert Opinions and Clinical Evidence

Leading foot and ankle surgeons widely regard Mann's surgery as a cornerstone in modern orthopedic practice. Multiple studies support its efficacy in correcting deformities, improving function, and maintaining joint integrity.

Key insights from experts include:

- The importance of comprehensive preoperative planning using advanced imaging techniques.
- The necessity of a multidisciplinary approach, involving radiologists, physical therapists, and rehabilitation specialists.
- Emphasis on patient education regarding realistic outcomes and postoperative care.

Recent research highlights:

- Improved patient satisfaction scores following minimally invasive Mann's procedures.
- Comparable or superior outcomes relative to traditional open surgeries.
- Reduced complication rates with meticulous surgical technique.

Patient Perspectives and Outcomes

Patients who undergo Mann's surgery often report:

- Significant pain relief
- Improved alignment and gait
- Preservation of joint mobility
- Faster return to daily activities and sports

However, success hinges on proper patient selection, adherence to postoperative protocols, and surgeon expertise.

Future Directions and Innovations

The landscape of foot and ankle surgery continues to evolve, with Mann's principles adapting to emerging technologies:

- 3D Printing and Custom Implants: Enhancing precision in osteotomies.
- Robotics and Navigation: Increasing surgical accuracy.
- Biologics and Regenerative Medicine: Supporting soft tissue and cartilage healing.
- Enhanced Rehabilitation Protocols: Accelerating recovery and functional restoration.

Conclusion: Is Mann's Surgery the Right Choice?

Mann's surgery of the foot and ankle represents a sophisticated, patient-centric approach that combines biomechanical precision with minimally invasive techniques. Its versatility makes it suitable for a wide range of deformities and degenerative conditions, offering the potential for durable, natural-function-preserving outcomes.

Key takeaways:

- It is best performed by surgeons with specialized training in foot and ankle biomechanics.
- Proper patient selection and comprehensive planning are critical.
- Ongoing research and technological advancements promise to further improve results.

For patients seeking to address complex foot and ankle issues with a procedure that respects natural anatomy and promotes quick recovery, Mann's surgery offers an expert-backed, innovative solution—an approach that continues to shape the future of orthopedic foot and ankle care.

Disclaimer: This review is intended for informational purposes and should not replace professional medical consultation. Always consult a qualified orthopedic surgeon or podiatrist to discuss individual conditions and treatment options.

Question What types of foot and ankle conditions does Mann's Surgery expert consultation typically address? Mann's Surgery expert consultation covers a wide range of conditions including bunions, hammertoes, heel spurs, ankle instability, Achilles tendinitis, and complex deformities requiring surgical intervention. **How can I prepare for my Mann's Surgery foot and ankle consultation?** Preparation involves bringing a detailed medical history, list of current medications, any previous imaging or reports, and being ready to discuss your symptoms, pain levels, and activity limitations with the specialist. **What are the common surgical procedures performed by Mann's Surgery experts for foot and ankle issues?** Common procedures include bunion correction (hallux valgus surgery), minimally invasive hammertoe repair, ankle stabilization surgeries, Achilles tendon repairs, and reconstructive surgeries for deformities. **What are the risks and recovery expectations after foot and ankle surgery with Mann's Surgery experts?** Risks may include infection, nerve injury, or delayed healing. Recovery typically involves a period of immobilization, physical therapy, and gradual return to activity, with guidance tailored to each procedure. **Are non-surgical treatments considered before opting for Mann's Surgery procedures?** Yes, conservative treatments like physical therapy, orthotics, medications, and injections are usually recommended first. Surgery is considered when these options fail to relieve symptoms or in cases of severe deformity. **How experienced are Mann's Surgery foot and ankle specialists in minimally invasive procedures?** Mann's Surgery experts are highly experienced in minimally invasive techniques, which often result in less pain, shorter recovery times, and better cosmetic outcomes for suitable cases. **Can Mann's Surgery experts customize treatment plans based on individual patient needs?** Absolutely. Treatment plans are personalized, considering the patient's specific condition, lifestyle, activity level, and health status to achieve optimal outcomes. **What should I expect during a consultation with Mann's Surgery foot and ankle specialist?** During the consultation, the specialist will evaluate your foot and ankle condition through physical examination and imaging if needed, discuss your symptoms, medical history, and outline possible treatment options. **How do I find the best Mann's Surgery foot and ankle expert for my condition?** Look for board-certified orthopedic surgeons with specialized training in foot and ankle surgery, check patient reviews, ask about their experience with your specific condition, and seek recommendations from your primary care provider. **What advancements or latest techniques are being used by Mann's Surgery experts in foot and**

ankle surgery? Experts utilize the latest techniques such as minimally invasive surgery, 3D preoperative planning, bio-absorbable implants, and arthroscopic procedures to improve precision, reduce recovery time, and enhance patient outcomes.

Related keywords: foot and ankle surgery, podiatric surgery, foot specialist, ankle surgery, orthopedic surgeon, foot and ankle expert, surgical consultation, minimally invasive foot surgery, sports injury foot surgery, foot and ankle treatment

Arthroscopy and endoscopy of the foot and ankle p

arthroscopy and endoscopy of the foot and ankle p

Arthroscopy and endoscopy of the foot and ankle represent minimally invasive surgical techniques that have revolutionized the management of various pathologies affecting these complex joints. These procedures utilize small incisions and specialized instruments to visualize, diagnose, and treat intra-articular and peri-articular structures with reduced morbidity compared to traditional open surgeries. As the foot and ankle are intricate anatomical regions comprising multiple bones, ligaments, tendons, cartilage, and neurovascular structures, these techniques demand specialized skills and thorough understanding of regional anatomy. Their growing popularity is driven by benefits such as decreased postoperative pain, faster recovery times, improved cosmetic outcomes, and the ability to address multiple pathologies simultaneously. This article offers a comprehensive exploration of arthroscopy and endoscopy in the foot and ankle, covering indications, surgical techniques, instrumentation, complications, and future perspectives.

Understanding Arthroscopy and Endoscopy in the Foot and Ankle

Definition and Principles

Arthroscopy involves the use of a fiberoptic camera (arthroscope) inserted into the joint through small portals to visualize intra-articular structures directly. Endoscopy, in the context of the foot and ankle, often refers to minimally invasive visualization of peri-articular regions, tendons, and other soft tissues outside the joint capsule. Both techniques rely on high-resolution imaging, specialized instruments, and minimally invasive access to perform diagnostic assessment and therapeutic interventions.

Key principles include:

- Use of small incisions (portals)
- Preservation of surrounding tissues
- Enhanced visualization of joint and peri-articular structures
- Precise instrumentation for debridement, repair, or removal of pathology

Historical Development

The advent of arthroscopy in the 20th century transformed joint surgery, initially in the knee and shoulder, before expanding to the ankle and foot. The first ankle arthroscopy was performed in the 1970s, with ongoing advances in instrumentation and technique. As understanding of foot and ankle pathologies evolved, so did the utilization of endoscopic procedures targeting specific regions such as the subtalar joint, talar dome, and tendon sheaths.

Indications for Arthroscopy and Endoscopy of the Foot and Ankle

Common Indications

The decision to employ arthroscopy or endoscopy is based on clinical evaluation, imaging, and patient-specific factors. Typical indications include:

- Osteochondral lesions of the talus
- Anterior and posterior ankle impingement syndromes
- Synovitis and joint inflammation
- Loose body removal
- Assessment and treatment of osteoarthritis
- Peroneal and tibialis posterior tendinopathy
- Haglund's deformity and retrocalcaneal bursitis
- Subtalar joint pathologies
- Evaluation and management of infections or cysts
- Correction of deformities such as varus or valgus deformities

Specific Conditions Managed

- Osteochondral Lesions of the Talus: Often caused by trauma, these lesions can lead to chronic ankle pain and instability. Arthroscopy allows for debridement and microfracture procedures.
- Ankle Impingement: Both anterior and posterior impingement syndromes benefit from arthroscopic debridement of osteophytes and soft tissue scar tissue.
- Peroneal Tendon Pathology: Tendon subluxation, tears, or tenosynovitis can be addressed via endoscopic tenosynovectomy or tendon repair.

- Subtalar and Midfoot Pathologies: Endoscopy allows visualization and treatment of sinus tarsi syndrome, cartilage lesions, and cysts.
- Haglund's Deformity: Endoscopic calcaneoplasty reduces postoperative morbidity compared to open procedures.

Instrumentation and Techniques

Equipment Used in Foot and Ankle Arthroscopy

- Arthroscopes: Ranging from 2.7mm to 4.0mm in diameter, with high-definition imaging capabilities.
- Light Sources and Cameras: Provide illumination and visualization.
- Trocars and Cannulas: For portal creation and instrument passage.
- Shavers and Debridement Tools: For tissue removal.
- Microfracture Needles: For cartilage repair.
- Suction and Irrigation Systems: To maintain clear visualization.
- Specialized Instruments: Including graspers, scissors, burrs, and radiofrequency devices.

Portal Placement Techniques

Proper portal placement is critical for access and safety:

- Anteromedial and Anterolateral Portals: Commonly used for anterior ankle procedures.
- Posterior Portals: For posterior ankle and subtalar joint access.
- Medial and Lateral Portals: For peroneal tendons and lateral gutter pathology.
- Sinus Tarsi Portal: For subtalar joint endoscopy.

Precise anatomical knowledge minimizes neurovascular injury risk and optimizes surgical outcomes.

Step-by-Step Surgical Approach

While specific techniques vary depending on pathology, a typical arthroscopic procedure involves:

- Patient Positioning: Supine or prone, with appropriate limb support.
- Landmark Identification: Using palpable landmarks and imaging guidance.
- Portal Creation: Under sterile conditions, small incisions are made, and cannulas are inserted.
- Joint or Region Visualization: The arthroscope is introduced, and intra-articular or peri-articular structures are inspected.

- Pathology Management: Debridement, removal of loose bodies, osteophyte resection, or repair as needed.
- Closure and Postoperative Care: Portals are closed, and postoperative protocols are initiated.

Complications and Limitations

Potential Complications

- Neurovascular injury, particularly to superficial nerves (superficial peroneal, sural nerve)
- Infection
- Hemarthrosis or bleeding
- Instrument breakage or iatrogenic cartilage damage
- Incomplete visualization or missed pathology
- Postoperative stiffness or pain

Limitations of Arthroscopy and Endoscopy

- Limited field of view compared to open surgery
- Steep learning curve requiring specialized training
- Contraindicated in cases of severe joint destruction or instability
- Not suitable for extensive deformities or complex trauma

Postoperative Management and Rehabilitation

Effective postoperative protocol enhances recovery:

- Immediate Postoperative Care: RICE (Rest, Ice, Compression, Elevation), pain control
- Weight-Bearing: Typically restricted initially; progression based on procedure
- Physical Therapy: Focused on range of motion, strength, and proprioception
- Monitoring: Regular follow-up with clinical and imaging assessments

Future Directions and Innovations

Advancements in technology continue to expand the scope and efficacy of arthroscopy and endoscopy in foot and ankle surgery:

- 3D Imaging and Navigation Systems: Improve accuracy and safety.

- Robotics: Potential for enhanced precision.
- Biological Augmentation: Use of stem cells and growth factors to promote healing.
- Miniaturization and Instrument Development: Allowing access to previously challenging areas.
- Enhanced Visualization: High-definition and 4K cameras facilitate better tissue assessment.

Conclusion

Arthroscopy and endoscopy of the foot and ankle are invaluable tools that offer minimally invasive options for diagnosing and treating a wide array of pathologies. Their success depends on meticulous surgical technique, comprehensive anatomical knowledge, and appropriate patient selection. As technology advances, these techniques will likely become even more precise and versatile, further improving patient outcomes. Surgeons must stay abreast of evolving methods and innovations to harness their full potential while minimizing risks. Ultimately, these procedures exemplify the shift toward less invasive, tissue-preserving approaches in orthopedic foot and ankle surgery, aligning with contemporary goals of rapid recovery and optimal functional restoration.

Arthroscopy and Endoscopy of the Foot and Ankle: A Comprehensive Review

The fields of arthroscopy and endoscopy of the foot and ankle have revolutionized the way orthopedic surgeons diagnose and treat a variety of musculoskeletal conditions affecting these complex regions. Minimally invasive techniques have replaced traditional open surgeries in many cases, offering patients reduced pain, shorter recovery times, and improved functional outcomes. This article provides an in-depth exploration of the principles, indications, procedures, advantages, limitations, and future prospects of arthroscopic and endoscopic interventions in the foot and ankle.

Introduction to Arthroscopy and Endoscopy of the Foot and Ankle

Arthroscopy and endoscopy are minimally invasive surgical techniques that utilize small incisions, specialized instruments, and a camera to visualize and treat internal structures within the joint or soft tissue compartments. Although often used interchangeably, arthroscopy generally refers to joint procedures, while endoscopy can encompass a broader range of soft tissue and bursal interventions.

In the foot and ankle, these techniques allow for precise treatment of various pathologies, including ligament injuries, cartilage lesions, Synovitis, impingements, and soft tissue cysts, with less morbidity compared to traditional open approaches.

Historical Development and Evolution

The advent of ankle arthroscopy dates back to the late 1970s, with initial techniques evolving rapidly through advances in instrumentation and imaging. Early procedures were limited by technical

difficulties and limited visualization, but innovations such as high-definition cameras, specialized portals, and improved instrumentation have expanded the scope of minimally invasive foot and ankle surgery.

Over time, the indications have broadened from diagnostic exploration to include complex therapeutic interventions, making arthroscopy and endoscopy essential tools in the modern foot and ankle surgeon's armamentarium.

Principles of Arthroscopy and Endoscopy in the Foot and Ankle

The core principles involve:

- Minimally invasive access: Small portals are created to introduce the camera and instruments.
- Visualization: High-definition cameras provide magnified views of intra-articular and peri-articular structures.
- Targeted intervention: Precise treatment minimizes collateral tissue damage.
- Reduced morbidity: Less tissue disruption leads to quicker recovery.

Understanding the anatomy, portal placement, and instrument handling is crucial for successful procedures.

Common Indications for Arthroscopy and Endoscopy

The indications can be broadly categorized into diagnostic and therapeutic:

Diagnostic indications:

- Unexplained ankle pain
- Persistent swelling
- Evaluation of intra-articular pathology

Therapeutic indications:

- Ankle impingement syndromes (anteromedial, posterior)
- Osteochondral lesions of the talus
- Synovitis and capsular adhesions
- Loose bodies and intra-articular foreign bodies
- Ligamentous injuries, including anterior talofibular ligament (ATFL) repairs

- Tarsal tunnel syndrome (endoscopic decompression)
- Soft tissue cysts, such as ganglia

Technical Aspects and Procedure Overview

Preoperative Planning

- Imaging studies (MRI, CT scans) help identify pathology and plan portals.
- Patient positioning (supine, lateral, or prone) depends on the target area.
- Adequate anesthesia and tourniquet application optimize visualization.

Portal Placement

- Standard portals include anteromedial, anterolateral, posterolateral, and posteromedial portals.
- Proper portal placement minimizes neurovascular injury risks.
- In some procedures, accessory portals are created for better access.

Instrumentation

- Arthroscopes with various diameters (2.7mm-4mm)
- Shavers, burrs, graspers, scissors, and laser devices
- Fluid management systems to maintain joint distension and clear visualization

Procedure Steps

- Establish portals under visualization
- Insert arthroscope and assess intra-articular structures
- Perform targeted interventions (e.g., debridement, microfracture, ligament repair)
- Confirm adequacy of treatment before closure

Specific Procedures in Foot and Ankle Arthroscopy

Ankle Arthroscopy

- Anterior ankle arthroscopy: common for anterior impingement, osteochondral lesions
- Posterior ankle arthroscopy: for posterior impingement, flexor hallucis longus (FHL)

tenosynovitis

- Lateral ankle arthroscopy: addressing ligamentous injuries and loose bodies

Subtalar and Hindfoot Arthroscopy

- Used for sinus tarsi syndrome, coalition resection, and osteochondral lesions

Midfoot and Forefoot Arthroscopy

- Corrects deformities like hallux valgus, addresses soft tissue pathologies

Endoscopic Techniques in Foot and Ankle

Endoscopy often pertains to soft tissue procedures, including:

- Tarsal Tunnel Endoscopy: for nerve decompression
- Extensor Retinaculum Endoscopy: for tendinopathy
- Bursal Endoscopy: for retrocalcaneal bursitis
- Ganglion cyst excision

These procedures utilize small incisions and specialized endoscopes to access soft tissue compartments with minimal disruption.

Advantages of Arthroscopy and Endoscopy

- Reduced postoperative pain and swelling
- Shorter hospital stay and faster mobilization
- Lower infection rates due to smaller incisions
- Enhanced visualization of intra-articular structures
- Improved cosmetic outcomes with minimal scarring
- Ability to perform both diagnostic and therapeutic procedures simultaneously

Limitations and Challenges

Despite the numerous benefits, certain limitations exist:

- Technical difficulty: Requires specialized training and experience
- Limited visualization in complex or scarred joints
- Risk of neurovascular injury if portals are misplaced
- Inadequate for large or complex deformities requiring open correction
- Steep learning curve for some procedures
- Cost of equipment and disposables

Complications and Risks

Potential complications include:

- Neurovascular injury
- Wound infection
- Hemarthrosis
- Instrument breakage
- Postoperative stiffness
- Incomplete treatment leading to persistent symptoms

Proper technique, thorough knowledge of anatomy, and careful patient selection are vital for minimizing risks.

Postoperative Care and Rehabilitation

Rehabilitation protocols vary depending on the procedure but generally include:

- Early mobilization within pain limits
- Weight-bearing as tolerated or protected
- Physical therapy focusing on range of motion, strength, and proprioception
- Regular follow-up to monitor healing and function

Future Directions and Innovations

Advances continue in:

- Navigation and imaging guidance: improving accuracy
- High-definition 3D visualization
- Robotics and computer-assisted techniques
- Biological therapies: such as stem cell and platelet-rich plasma (PRP) injections combined with

minimally invasive procedures

- Enhanced instrumentation for better access to hard-to-reach areas

These innovations promise to expand indications and improve outcomes further.

Conclusion

Arthroscopy and endoscopy of the foot and ankle represent a significant leap forward in orthopedic surgery, offering minimally invasive options for diagnosing and treating a wide array of pathologies. Their success hinges on meticulous technique, thorough understanding of anatomy, and appropriate case selection. While they are not suitable for all conditions, their role continues to grow, driven by technological advancements and clinical evidence supporting improved patient outcomes. As expertise and equipment evolve, the scope of arthroscopic and endoscopic interventions will undoubtedly expand, marking an exciting future for foot and ankle surgery.

In summary, the integration of arthroscopy and endoscopy into foot and ankle surgery has transformed patient care by enabling precise, less invasive interventions. Continued research and innovation are essential to overcome current limitations and fully realize their potential.

Question What are the common indications for foot and ankle arthroscopy and endoscopy? Common indications include synovitis, loose body removal, osteoarthritis management, ligament repair, and evaluation of persistent pain or instability not responsive to conservative treatment. **Answer** How does arthroscopy of the foot and ankle differ from traditional open surgery? Arthroscopy involves minimally invasive techniques using small incisions and a camera, resulting in less tissue trauma, faster recovery, and reduced postoperative pain compared to open surgery. What are the potential risks and complications associated with foot and ankle arthroscopy? Risks include nerve or vascular injury, infection, blood clot formation, joint stiffness, incomplete symptom relief, and, rarely, damage to surrounding structures. How long is the typical recovery period after ankle arthroscopy? Recovery varies depending on the procedure but generally ranges from a few days to several weeks, with most patients resuming light activities within 2-4 weeks and full recovery in 6-12 weeks. What advancements have recently been made in endoscopic techniques for foot and ankle surgery? Recent advancements include high-definition imaging, improved instrumentation, and the use of smaller portals, which enhance visualization, reduce complications, and expand the scope of minimally invasive procedures. Which conditions of the foot and ankle are best treated with arthroscopy? Conditions such as anterior ankle impingement, osteochondral lesions, synovitis, loose bodies, and certain ligament injuries are effectively treated with arthroscopy. Are there any contraindications for performing foot and ankle arthroscopy? Contraindications include active infection, significant joint deformity requiring open surgery, severe osteoarthritis, and patient comorbidities that impair healing or anesthesia risk. What are the benefits of using endoscopy for treating posterior ankle pathology? Endoscopy allows for minimally invasive access to posterior ankle structures, providing pain relief, reducing tissue disruption, and facilitating quicker recovery.

compared to open procedures.

Related keywords: foot arthroscopy, ankle endoscopy, minimally invasive foot surgery, ankle arthroscopy, foot joint surgery, ankle ligament repair, subtalar arthroscopy, ankle synovectomy, osteoarthritis foot, ankle cartilage repair

Release mcl at mtp joint cpt

Release MCL at MTP Joint CPT: A Comprehensive Guide to the Procedure, Indications, and Coding

Understanding the intricacies of musculoskeletal procedures is essential for healthcare providers, coders, and patients alike. One such procedure gaining attention is the release of the medial collateral ligament (MCL) at the metatarsophalangeal (MTP) joint, which involves specific surgical techniques and precise coding for reimbursement. This article delves into the details of performing the release of the MCL at the MTP joint, covering the indications, surgical techniques, postoperative considerations, and the appropriate CPT coding.

What is the Release of MCL at the MTP Joint?

The medial collateral ligament (MCL) of the MTP joint is a key stabilizer located on the medial (inner) side of the joint connecting the proximal phalanx to the metatarsal bone. In certain pathological conditions?such as severe hallux valgus deformity, recurrent bunions, or joint contractures?releasing or releasing and reconstructing the MCL may be necessary to realign the toe or restore joint function.

The release of the MCL at the MTP joint involves surgically cutting or lengthening the ligament to correct deformity or improve joint mobility. This is often performed in conjunction with other procedures like osteotomies or soft tissue balancing.

Indications for MCL Release at the MTP Joint

Understanding when to perform an MCL release is critical. Typical indications include:

- Severe hallux valgus deformity unresponsive to conservative management
- Recurrent bunion deformities after previous surgeries
- Rigid or flexible deformities requiring soft tissue balancing

- Joint contractures limiting toe movement
- Corrective procedures in congenital or acquired deformities

Proper patient selection ensures optimal outcomes and minimizes complications.

Surgical Technique for MCL Release at the MTP Joint

Performing an MCL release demands meticulous technique to preserve joint stability and prevent over-release. The general steps include:

Preparation

- The patient is positioned supine with a tourniquet applied to the thigh.
- Anesthesia is administered, typically regional or general anesthesia.
- The surgical site is prepared and draped in sterile fashion.

Approach

- A dorsal or medial incision is made over the MTP joint, depending on the surgeon's preference.
- Soft tissues are carefully dissected to expose the joint capsule and collateral ligament.

Identification of the MCL

- The MCL is identified as a thick, fibrous band on the medial side of the joint.
- The surgeon assesses the extent of tightness or deformity necessitating release.

Ligament Release

- The MCL is incised longitudinally or partially released, taking care to preserve surrounding structures.
- In some cases, a Z-plasty or lengthening technique may be employed to avoid complete detachment.
- The surgeon verifies the correction of deformity and joint stability intraoperatively.

Closure

- The capsule and soft tissues are repaired or balanced as needed.
- Skin is closed in layers, and a sterile dressing is applied.

Postoperative Care and Rehabilitation

Post-surgical management is crucial for optimal recovery:

- Immobilization with a postoperative shoe or boot for 2-6 weeks
- Elevation and ice to reduce swelling
- Gradual range-of-motion exercises as tolerated
- Weight-bearing as tolerated or restricted based on concomitant procedures
- Follow-up visits to monitor healing and address complications

Rehabilitation protocols vary depending on the extent of the release and associated procedures.

Understanding CPT Codes for MCL Release at MTP Joint

Accurate coding is essential for proper reimbursement. The Current Procedural Terminology (CPT) provides specific codes for soft tissue procedures of the foot.

Relevant CPT Codes

- 28122: Tenotomy, lengthening or release of tight or contracted tendons, fasciotomy, or soft tissue procedures around toe joints (e.g., for hammer toe, claw toe, or bunion correction)
- 28285: Osteotomy, bunionectomy (e.g., Austin, chevron, or akin osteotomy), with or without soft tissue correction
- 28705: Arthrodesis, toe joint (e.g., MTP, PIP, DIP), with or without fusion, with or without internal fixation

While there isn't a CPT code explicitly titled "release of MCL at MTP joint," CPT 28122 often encompasses soft tissue releases around toe joints, including ligament releases, when performed as part of deformity correction.

CPT Coding Tips:

- When the procedure involves only the ligament release without osteotomy or fusion, CPT 28122 is typically appropriate.
- If the release is performed in conjunction with other procedures such as osteotomy or joint fusion, modifiers (e.g., -51 for multiple procedures) and additional codes should be used.
- Document the procedure thoroughly, including the specific ligament released, to support the code selection.

Documentation Considerations

Proper documentation should include:

- Indication for the procedure
- Description of the ligament released and the technique used
- Details of any additional procedures performed
- Intraoperative findings
- Postoperative plan

Clear documentation ensures accurate coding and billing.

Common Complications and Their Management

Possible complications of MCL release include:

- Over-release leading to joint instability
- Infection
- Hematoma formation
- Persistent deformity
- Neurovascular injury

Preventive measures involve meticulous surgical technique, thorough knowledge of anatomy, and postoperative monitoring.

Conclusion

The release of the MCL at the MTP joint is a valuable procedure in the corrective arsenal for deformities such as hallux valgus. Its success depends on proper patient selection, precise surgical technique, and accurate CPT coding for reimbursement. Healthcare providers should stay familiar with the relevant codes, such as CPT 28122, and ensure comprehensive documentation to facilitate smooth billing processes.

By understanding the indications, techniques, and coding nuances, surgeons and coders can optimize patient outcomes and ensure compliance with coding standards. As foot and ankle surgery advances, staying current with procedural codes and best practices remains essential for delivering high-quality care.

Release MCL at MTP Joint CPT: An In-Depth Review and Guide

The release MCL at MTP joint CPT (Current Procedural Terminology) code refers to a specific surgical intervention aimed at releasing the medial collateral ligament (MCL) at the metatarsophalangeal (MTP) joint. This procedure is often performed to address contractures, deformities, or persistent pain caused by MCL tightness or fibrosis, especially in conditions like hallux valgus or other deformities affecting the great toe. Understanding the nuances of this procedure, including its indications, techniques, coding, and outcomes, is crucial for orthopedic surgeons, podiatrists, and coding professionals.

Understanding the Anatomy and Indications for MCL Release at MTP Joint

Anatomy of the MCL at the MTP Joint

The medial collateral ligament (MCL) at the MTP joint is a vital structure providing medial stability to the toe, especially the great toe. It spans from the medial condyle of the metatarsal head to the proximal phalanx, helping maintain proper alignment and prevent valgus deformity. The MCL is composed of superficial and deep fibers, making it a robust ligament that can sometimes become pathologically tight or fibrotic.

Indications for MCL Release

The primary indications for releasing the MCL at the MTP joint include:

- Persistent contracture or tightness causing deformity or functional impairment.
- Hallux valgus deformity where medial soft tissue release, including the MCL, is part of corrective procedures.
- Recurrent deformities after previous surgical correction.
- Pain localized to the medial aspect of the MTP joint unresponsive to conservative management.
- Limited toe motion impeding normal gait or footwear fitting.

It is crucial to evaluate whether the MCL tightness is contributing to the deformity and if releasing it will restore proper alignment without compromising joint stability.

Techniques for Releasing the MCL at MTP Joint

Surgical Approaches

There are various surgical techniques to release the MCL, tailored to the specific pathology and deformity severity:

- Open Release: A standard dorsal or medial approach allows direct visualization and precise release of the MCL fibers.
- Percutaneous Release: Minimally invasive technique involving small incisions and blunt dissection, suitable for less severe contractures.
- Z-Plasty or Lengthening Techniques: For cases where a simple release may weaken the joint, lengthening approaches may be employed.

Step-by-Step Overview of Open MCL Release

- Anesthesia and Positioning: Usually performed under local anesthesia with sedation or regional block, with the patient in supine position.
- Incision: A medial incision over the MTP joint is made, carefully avoiding neurovascular structures.
- Dissection: Soft tissues are gently dissected to expose the MCL.
- Ligament Release: The ligament fibers are carefully incised or divided, ensuring preservation of surrounding tissues.
- Assessment: Range of motion and deformity correction are evaluated intraoperatively.
- Closure and Postoperative Care: Incision is closed in layers, and postoperative protocols are initiated.

The choice of technique depends on the degree of deformity, soft tissue quality, and surgeon preference.

Understanding CPT Coding for MCL Release at MTP Joint

Relevant CPT Codes

The CPT code most associated with the release of the MCL at the MTP joint is:

- 28010 ? Release of joint capsule, toe, with or without removal of bursa and/or synovium; first toe (e.g., for deformity correction)

However, more specific codes may apply depending on the extent of the procedure:

- 28015 ? Surgical correction of hallux valgus (bunion), with or without sesamoidectomy; with soft tissue release and/or osteotomy

In some cases, the procedure might be bundled or combined with other interventions, such as osteotomies or arthrodesis, which require different codes.

Modifiers and Documentation

Proper documentation is critical to substantiate the CPT coding:

- Clearly note the indication for the release.
- Describe the extent of the ligament release.
- Specify whether the procedure was open or percutaneous.
- Document any additional procedures performed simultaneously.

Modifiers such as 51 (multiple procedures) or 59 (distinct procedural service) may be applicable depending on the scenario.

Pros and Cons of MCL Release at the MTP Joint

Pros:

- Deformity Correction: Allows realignment of the toe, especially in hallux valgus cases.
- Improved Functionality: Restores toe motion and gait mechanics.
- Minimally Invasive Options: Percutaneous techniques reduce recovery time and scarring.
- Adjunct to Other Procedures: Complements osteotomies or arthrodesis for comprehensive correction.

Cons:

- Risk of Over-release: Excessive release may lead to joint instability or medial subluxation.
- Neurovascular Injury: Potential damage to medial digital nerves or arteries.
- Recurrence: Contracture or deformity may recur if soft tissue balancing is inadequate.
- Postoperative Stiffness: Scar formation could limit toe mobility if not managed properly.

Postoperative Management and Outcomes

Postoperative Care

- Immobilization: Usually involves a stiff-soled shoe or postoperative boot.
- Weight-bearing: Often permitted as tolerated, depending on concomitant procedures.
- Physical Therapy: Emphasized to regain range of motion and strength.
- Monitoring: Regular follow-up to detect signs of instability, infection, or recurrence.

Expected Outcomes

Most patients experience:

- Significant improvement in toe alignment.
- Enhanced toe mobility.
- Reduction in medial pain.
- Better footwear fit and gait pattern.

However, outcomes depend on the severity of initial deformity, surgical technique, and adherence to postoperative protocols.

Conclusion

The release MCL at MTP joint CPT encompasses a critical surgical step in correcting toe deformities, especially hallux valgus. Proper understanding of the anatomy, indications, surgical techniques, and coding nuances ensures optimal patient outcomes and accurate billing. While the procedure offers significant benefits in deformity correction and functional improvement, surgeons must weigh the potential risks and ensure meticulous technique. As with all surgical interventions, comprehensive patient evaluation, careful planning, and adherence to postoperative management are paramount for success.

In summary:

- The procedure is indicated mainly for deformity correction and contracture release.
- Multiple surgical approaches exist, each with its indications.
- Accurate CPT coding requires detailed operative documentation and understanding of the procedure extent.
- The benefits often outweigh the risks when performed judiciously.

- Proper postoperative care is essential for sustained results.

By mastering the intricacies of the release MCL at MTP joint CPT, clinicians can improve patient outcomes and ensure compliance with coding standards, ultimately enhancing the quality of foot and ankle care.

Question What does the CPT code for releasing the MCP joint involve? The CPT code for releasing the MCP joint typically involves codes 26520 or 26521, which cover surgical procedures to release contractures or adhesions at the metacarpophalangeal joint, depending on the complexity of the procedure. When is a release of the MCP joint at the MTP joint indicated? A release of the MCP joint at the MTP joint is indicated in cases of severe contracture, stiffness, or deformity that impair hand function, often due to conditions like rheumatoid arthritis or trauma. What are the key considerations for coding MCP joint release procedures in CPT? Key considerations include accurately documenting the specific procedure performed, whether it's a simple release or an extensive release, and ensuring the correct CPT code (such as 26520 or 26521) is used based on the procedure's complexity. Are there any modifiers required when reporting MCP joint release procedures at the MTP joint? Modifiers may be necessary if the procedure is bilateral (modifier 50), performed as an add-on, or if multiple procedures are done during the same session. Always review payer-specific guidelines for accurate modifier use. What are the post-operative considerations following MCP joint release at the MTP joint? Post-operative care typically includes immobilization, physical therapy to restore range of motion, and monitoring for complications such as infection or joint instability to ensure optimal recovery. How do the CPT codes differ between simple and complex MCP joint releases? Simple MCP joint release procedures are usually coded as 26520, while more complex or extensive releases involving additional procedures or reconstructions may be coded as 26521, reflecting the increased complexity. What documentation is essential to support CPT coding for MCP joint release at the MTP joint? Essential documentation includes detailed operative reports describing the procedure performed, the indication for surgery, the extent of release, and any additional procedures performed to justify the selected CPT code.

Related keywords: release mcl at mtp joint, mcp joint ligament release, metatarsophalangeal joint surgery, mcl release procedure, CPT code for mcl release, mtp joint ligament procedure, metatarsal phalangeal joint surgery, mcl release CPT, surgical release of mcl at mtp, mcl repair at mtp joint